

Pre-operative differentiation of atypical parathyroid tumors and parathyroid adenomas: is it possible?

¿Es posible diferenciar preoperatoriamente los tumores paratiroides atípicos de los adenomas paratiroides?

Ibrahim Kılınc^{1*}, Serap Ulusoy¹, Mustafa Oruç¹, Alparslan Ertenlice¹, Aydan Kılıçarslan², and Tolga Dinç¹

¹Department of General Surgery; ²Department of Pathology. Ankara Bilkent City Hospital, Ankara, Türkiye

Abstract

Objectives: This study aimed to determine whether atypical parathyroid tumors (APTs) can be differentiated from benign parathyroid adenomas (PAs) preoperatively and to compare their biochemical characteristics, post-operative complications, and long-term outcomes. **Methods:** A retrospective analysis was conducted on patients who underwent parathyroidectomy between 2010 and 2022. Fifty patients with APT and one hundred with PA (control group) were included. Demographic data, biochemical parameters, complications, and follow-up outcomes were compared. **Results:** Pre-operative serum calcium and parathyroid hormone (PTH) levels were significantly higher in APT patients. Receiver operating characteristic analysis identified $\text{Ca} > 11.5 \text{ mg/dL}$ and $\text{PTH} > 180 \text{ pg/mL}$ as thresholds for distinguishing APTs. Multivariate analysis confirmed these as independent predictors. APT patients had a higher rate of concurrent thyroid surgery and more frequent complications such as transient nerve injury and hungry bone syndrome. No significant difference was found in persistent disease or recurrence after an average follow-up of 4.8 years. **Conclusions:** Pre-operative Ca and PTH levels may help distinguish APTs from PAs. Patients with APT should be informed about the increased risk of complications and concurrent thyroid surgery, and appropriate preventive strategies should be considered.

Keywords: Atypical parathyroid tumor. Parathyroid adenoma. Calcium. Parathyroid hormone nerve paralysis. Hungry bone syndrome.

Resumen

Objetivos: Determinar si los tumores paratiroides atípicos (TPA) pueden diferenciarse de los adenomas paratiroides (AP) en el periodo preoperatorio, y comparar sus características bioquímicas, complicaciones posoperatorias y evolución a largo plazo. **Métodos:** Se realizó un análisis retrospectivo de pacientes sometidos a paratiroidectomía entre 2010 y 2022. Se incluyeron 50 pacientes con TPA y 100 con AP como grupo control. Se compararon datos demográficos, parámetros bioquímicos, complicaciones quirúrgicas y resultados del seguimiento. **Resultados:** Los niveles preoperatorios de calcio sérico y hormona paratiroidea (PTH) fueron significativamente más elevados en los pacientes con TPA. El análisis ROC identificó calcio $> 11.5 \text{ mg/dL}$ y $\text{PTH} > 180 \text{ pg/mL}$ como puntos de corte útiles para distinguir los TPA. El análisis multivariado confirmó estos valores como predictores independientes. Los pacientes con TPA presentaron mayor frecuencia de cirugía tiroidea concomitante y complicaciones como lesión nerviosa transitoria y síndrome del hueso hambriento. No se observaron diferencias significativas en enfermedad persistente ni en recurrencia tras un seguimiento promedio de 4.8 años. **Conclusiones:** Los niveles

*Correspondence:

Ibrahim Kılınc

E-mail: ikilinc8083@gmail.com

Date of reception: 24-04-2025

Date of acceptance: 05-09-2025

DOI: 10.24875/CIRU.25000243

Cir Cir. 2026;94(4):422-430

Contents available at PubMed

www.cirugiaycirujanos.com

0009-7411/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

elevados de calcio y PTH preoperatorios pueden contribuir al diagnóstico diferencial de TPA. Se debe informar a los pacientes sobre el mayor riesgo quirúrgico y considerar estrategias preventivas.

Palabras clave: Tumor paratiroideo atípico. Adenoma paratiroideo. Calcio. Hormona paratiroidea. Parálisis nerviosa. Síndrome del hueso hambriento.

Introduction

Parathyroid tumors are a relatively homogeneous group of neoplasms, affecting approximately 0.1-0.3% of the population. These tumors are primarily characterized by the dysregulation in the secretion of parathyroid hormone (PTH), resulting in primary hyperparathyroidism (PHPT).¹ Among patients with PHPT, around 10% present with parathyroid hyperplasia, 85% with a single parathyroid adenoma (PA), and 3% with multiple PAs, whereas fewer than 2% of cases involve atypical parathyroid tumors (APT) or parathyroid carcinomas (PC).²

APTs represent a distinct clinical entity, a state considered to be in between benign PAs and malignant PCs. These tumors exhibit atypical cytological and/or architectural characteristics but do not demonstrate definitive evidence of capsular, vascular, perineural, or adjacent tissue invasion, nor metastasis. Previously termed atypical PA, the nomenclature has been updated to APT in the 5th edition of the World Health Organization (WHO) classification of endocrine tumors.³

The classification of APTs has been controversial due to their ambiguous features, both clinically and histopathologically. However, in 2018, the American Joint Committee on Cancer added the “Tis” category to the staging of parathyroid cancers, defining “Tis” as an atypical parathyroid neoplasm with uncertain malignant potential.⁴ Similarly, according to the 2022 WHO classification, APTs are categorized as neoplasms of uncertain malignant potential.

Symptoms and findings in APT patients include nephrolithiasis, fatigue, bone pain, brown tumors, and depression. Importantly, these findings are not specific to APTs.² Most APTs behave similarly to PAs and do not recur after surgical excision. However, recurrent disease and metastasis have been reported, particularly in cases with parafibromin (PFIB) (a tumor suppressor protein, PFIB) deficiency.⁵⁻⁸ Therefore, management of APTs is recommended to consist solely of clinical and biochemical follow-up.³

The primary aim of this study was to identify distinct morphological and laboratory characteristics of APTs,

beyond histopathological findings, that differentiate them from PAs and to establish pre-operative markers suggestive of APT. The secondary aim was to compare the short- and long-term outcomes of these two parathyroid neoplasms.

Methods

In this retrospective study, data were collected from patients who underwent surgery for PHPT at our hospital between January 2010 and December 2022. A total of 486 patients underwent surgery during this period. Of these, 77 patients were excluded due to chronic renal failure (n = 12), secondary or tertiary hyperparathyroidism (n = 9), familial syndromes, including MEN1 and MEN2A (n = 4), carcinoma (n = 6), and incomplete medical records (n = 46). Ultimately, 359 patients were diagnosed with PA, and 50 with APT. Fifty patients diagnosed with APT and one hundred patients with PA, who were treated during the same period, were randomly selected for inclusion in the study. To minimize selection bias, the 100 PA patients were selected from the 351 eligible cases using R statistical software. Each patient was assigned a unique identifier, and simple random sampling without replacement was performed with a fixed random seed to ensure reproducibility and maintain methodological rigor.

Demographic data (age, sex), date of surgery, date of the last evaluation (the most recent serum calcium [Ca] and PTH levels and/or neck imaging recorded in the medical files were considered as the date of the last evaluation), concurrent thyroid surgery was documented. Tumor localization, size, volume (parathyroid tumors were considered as ellipsoid objects, and the volume was calculated using the formula $\frac{4}{3} \pi abc$ (where a, b, and c represent the radii)). Biochemical data included pre-operative serum calcium (Ca, mg/dL) and PTH (pg/mL) levels, which were obtained from the records. For the pre-operative values, the highest measurements within the month preceding surgery were considered. The reference ranges for Ca and PTH were 8.5-10.5 mg/dL and 15-60 pg/mL, respectively. Histopathological features observed in APTs

were documented. Microscopy findings in pathology reports were re-evaluated based on the 2022 WHO classification updates for parathyroid tumors and ranked according to their frequency.

Recurrent laryngeal nerve injury was defined as a known injury occurring during surgery, as documented in the operative report, and/or based on evidence of hypomobility or immobility observed upon post-operative clinical laryngoscopy. No improvement in nerve function within 6 months was considered as permanent nerve damage. The absence of a decrease in PTH and Ca levels in the post-operative period was considered as persistent disease. Disease recurrence was defined as the reappearance of hypercalcemia along with elevated serum PTH levels after a disease-free period (normocalcemia) of at least 6 months. Patients who, within the first seven post-operative days and in the presence of normal or elevated PTH levels, developed hypocalcemia symptoms (e.g., paresthesia or muscle spasms) requiring intravenous Ca supplementation, or exhibited a serum Ca concentration below 7.5 mg/dL, were considered to have hungry bone syndrome (HBS).

This study was approved by the Ethics Committee of Ankara Bilkent City Hospital (Approval number E2-23-3377).

Statistical analysis

A power analysis was conducted to determine the required sample size for detecting a significant difference between PA and atypical parathyroid adenoma (APT) groups. The effect size was set at 0.50 (moderate effect), with an alpha level of 0.05 and a power of 80% ($\beta = 0.20$). Due to the unequal sample allocation ($n_2/n_1 = 0.5$), the required total sample size was adjusted accordingly. Using G*Power 3.1, the estimated minimum sample size for an independent samples t-test with this allocation ratio was $n_1 = 96$ (PA), $n_2 = 48$ (APT), total = 132. For this reason, this study included 100 PA and 50 APT patients.

All statistical analyses were performed using R (version 4.4.1). Continuous variables were tested for normality using the Shapiro–Wilk test. Normally distributed data were presented as mean $\pm$ standard deviation and compared using the independent samples t-test. Non-normally distributed data were expressed as median and interquartile range (IQR, 25–75th percentile) and compared using the Mann-Whitney U test. Categorical variables were presented as frequencies and percentages (%) and compared using the

Chi-square test or Fisher's exact test, as appropriate. A receiver operating characteristic (ROC) curve analysis was performed to determine the diagnostic performance of pre-operative calcium and PTH levels in distinguishing atypical parathyroid adenoma (APT) from PA. The area under the curve (AUC) was calculated, and optimal cutoff values were determined. Sensitivity, specificity, positive predictive value, and negative predictive value were reported. To identify independent predictors of APT, a multivariate logistic regression analysis was performed, including variables that were statistically significant in univariate analysis ($p < 0.05$). Results were expressed as odds ratios (OR) with 95% confidence intervals (CI). The goodness-of-fit of the final model was assessed using the Hosmer-Lemeshow test.

Results

A total of 150 patients were included in the study, with 100 diagnosed with PA and 50 with APT. The mean age of patients with PA was significantly higher (56 ± 13 years) compared to those with APT (50 ± 12 years, $p < 0.005$). The sex distribution did not differ significantly between the two groups ($p = 0.899$).

The median lesion diameter was larger in APT cases (20 mm, IQR 15–27) compared to PA (17 mm, IQR 13–21), though this difference did not reach statistical significance ($p = 0.054$). The median lesion volume was also higher in APT patients (996.5 mm^3 , IQR 563–2612) compared to PA (734.34 mm^3 , IQR 370.44–1382.3), but the difference was not statistically significant ($p = 0.085$) as shown in table 1. The distribution of lesion localization differed significantly between groups ($p < 0.05$). APT lesions were most frequently found in the left superior and right inferior positions, while PA lesions showed a more even distribution. Concomitant thyroid surgery was significantly more common in APT patients, with 42% undergoing lobectomy and 32% requiring total thyroidectomy, compared to 20% and 12%, respectively, in PA patients ($p < 0.005$).

Preoperatively, APT patients exhibited significantly higher PTH and Ca levels compared to PA patients. The median pre-operative PTH level in APT cases was 242.75 pg/dL (IQR 185–407.5), significantly higher than in PA cases (166.3 pg/dL , IQR 127.5–268.2; $p < 0.001$). Similarly, pre-operative calcium levels were significantly elevated in APT patients (11.93 mg/dL , IQR 11.33–12.2) compared to PA patients (11.03 mg/dL , IQR 10.79–11.72; $p < 0.001$). However,

Table 1. Patients demographics, adenoma characteristics, and laboratory measurements

Variables	Parathyroid adenoma (n = 100)	Atypical parathyroid tumor (n = 50)	p
Age*	56 ± 13	50 ± 12	< 0.005
Sex†	29 (29)	15 (30)	0.899
Median lesion diameter (mm)‡	17 (13-21)	20 (15-27)	0.054
Volume (mm³)‡	734.34 (370.44-1382.3)	996.5 (563-2612)	0.085
Localization†			
Left superior	9 (9)	10 (20)	< 0.05
Left inferior	45 (45)	12 (24)	
Right inferior	34 (45)	20 (40)	
Right superior	12 (12)	8 (16)	
Concurrent thyroid surgery†			
None	68 (68)	21 (42)	< 0.005
Lobectomy	12 (12)	16 (32)	
Thyroidectomy	20 (20)	13 (26)	
Pre-operative Pth (pg/dL)‡	166.3 (127.5-268.2)	242.75 (185-407.5)	< 0.001
Pre-operative Ca (mg/dL)‡	11.03 (10.79-11.72)	11.93 (11.33-12.2)	< 0.001
Post-operative PTH (pg/dL)‡	23 (13-36)	30.15 (12.52-52.10)	0.369
Post-operative Ca (mg/dL)‡	9 (8.6-9.39)	8.95 (8.51-9.32)	0.354
Pth > 180 pg/mL‡	42 (42)	38 (76)	< 0.001
Ca > 11.5 mg/dL‡	31 (31)	35 (70)	< 0.001
Last PTH measurement (pg/dL)‡	44.85 (35.20-54.81)	44.35 (34.9-56.3)	0.897
Last Ca measurement (mg/dL)‡	9.50 (8.44-9.52)	9.40 (8.90-9.62)	0.444

*Values presented as the mean ± standard deviation.

†Values presented as the number of patients (percentage).

‡Values presented as the median days (25-75th percentile).

post-operative PTH and Ca levels were not significantly different between the groups ($p = 0.369$ and $p = 0.354$, respectively).

Post-operative complications were more frequent in APT patients. Temporary recurrent nerve paralysis occurred in 16% of APT cases versus 6% of PA cases ($p < 0.05$). The incidence of “hungry bone syndrome” was also significantly higher in APT patients (16% vs. 3%, $p < 0.005$). Permanent recurrent nerve paralysis, post-operative disease persistence, and recurrence rates did not significantly differ between groups as shown in table 2.

Table 3 demonstrates the frequency of histopathological features observed in APT cases. Among APT cases, fibrotic bands were the most frequently observed histopathological feature (84%), followed by tumor cells in the capsule (40%) and atypical mitosis (36%). Trabecular growth was present in 28% of cases, while adhesion and Ki-67 proliferation index > 5% were each observed in 18% of cases. Increased

mitotic activity (> 5/10 mm²) and coagulative necrosis were the least common feature (4%) (Fig. 1).

Table 4 provides the details of the ROC curve analysis, and figures 2 and 3 illustrate the ROC curves for pre-operative PTH and Ca, respectively. ROC analysis was performed to determine the optimal cutoff values for distinguishing APT from PA. The AUC for calcium was 0.751 (95% CI: 0.672-0.830, $p < 0.001$), with an optimal cutoff of 11.5 mg/dL, yielding a sensitivity of 70% and specificity of 69%. The AUC for PTH was 0.691 (95% CI: 0.607-0.775, $p < 0.001$), with an optimal cutoff of 180 pg/dL, demonstrating a sensitivity of 76% and specificity of 58%. According to these results, A significantly higher proportion of APT patients had PTH levels exceeding 180 pg/mL (76% vs. 42%, $p < 0.001$) and calcium levels > 11.5 mg/dL (70% vs. 31%, $p < 0.001$).

Multivariate logistic regression analysis identified calcium > 11.5 mg/dL as an independent predictor of APT (OR: 3.571, 95% CI: 1.604-7.950, $p < 0.005$),




Figure 1. Histological features of atypical parathyroid tumor. Histological features of atypical parathyroid tumor. **A:** solid growth pattern formed by oncocytic cells and the thick fibrous capsule that develops it (x100 hematoxylin and eosin [H&E]). **B and C:** extension into the capsule that does not completely exceed the characteristics of the chief cell (x40 H&E). **D:** band-like fibrosis within the tumor (x100 H&E). **E:** increased mitotic figure, in square (x400 H&E). **F:** trabecular growth pattern and pleomorphism (x200 H&E). **G:** coagulative necrosis within the tumor (x200 H&E). **H:** growth staining in Ki-67 immunohistochemical test (x200).

Table 2. Post-operative complications and follow-up results

Complications	Parathyroid adenoma (n = 100)	Atypical parathyroid tumor (n = 50)	p
Temporary recurrent nerve paralysis†	6 (6)	8 (16)	< 0.05
Permanent recurrent nerve paralysis†	1 (1)	0 (0)	0.478
Hungry bone syndrome†	3 (3)	8 (16)	< 0.005
Temporary hypoparathyroidy†	5 (5)	4 (8)	0.466
Permanent hypoparathyroidy†	1 (1)	1 (1)	0.615
Persistent disease	3	2	
Recurrence disease	0	0	

*Values presented as the mean ± standard deviation.
†Values presented as the number of patients (percentage).

Table 3. Histopathological features of atypical parathyroid tumors

Histopathological features	Frequency n (%)
Band-like fibrosis	42 (84)
Tumor cells in capsule	20 (40)
Atypical mitotic figures	18 (36)
Trabecular growth	14 (28)
Adherence to adjacent structures	9 (18)
Ki-67 labeling index > 5%	9 (18)
Increased mitotic activity (> 5/10 mm²)	2 (4)
Coagulative necrosis	2 (4)

Discussion

In this study, our primary aim was to identify distinct morphological and laboratory characteristics of APTs, beyond histopathological findings, that differentiate them from PAs and to establish pre-operative markers suggestive of APT. Our secondary aim was to compare the post-operative complications and long-term follow-up outcomes of these two parathyroid neoplasms.

along with PTH > 180 pg/dL (OR: 1.127, 95% CI: 1.127-6.069, p < 0.05). Tumor volume did not show a significant association (p = 0.886). Table 5 presents the results of the multivariate analysis.

Table 4. Areas under curve for biochemical parameters for distinguishing PA from APT

Variables	AUC	95% CI	p	Cut-off	Sensitivity	Specificity	PPV	NPV
Ca	0.751	0.672-0.830	< 0.001	> 11.48	0.70	0.69	0.53	0.82
PTH	0.691	0.607-0.775	< 0.001	> 182	0.76	0.58	0.48	0.83

AUC: area under the curve; CI: confidence interval; PA: parathyroid adenoma; APT: atypical parathyroid adenoma; PPV: positive predictive value; NPV: negative predictive value; PTH: parathyroid hormone.

Table 5. Multivariate analysis of variables independently associated with the presence of APTs

Variables	Odds ratio	95% CI		Wald	p
		Lower	Upper		
Ca > 11.5 mg/dL	3.571	1.604	7.950	9.716	< 0.005
Pth > 180 pg/dL	1.127	1.127	6.069	5.007	< 0.05
Volume	1.000	-	-	0.021	0.886

Consistent with the literature,^{9,10} no significant difference in gender distribution was observed between patients with APT and PA. However, both APTs and PAs were more frequently diagnosed in females. Although some previous studies suggest that APTs and PAs typically occur at similar ages,^{9,11} the present study found that APTs manifested at a younger age (median age: 50 ± 12) than PAs. Indeed, Chen et al. evaluated 79 APT patients and reported that APTs were diagnosed at a younger age (median age: 46.9 ± 17.1) than PAs, attributing this to possible germline mutations.¹²

Many studies have compared sizes of parathyroid neoplasms. Although Marini et al. found no significant size difference between atypical tumors and carcinomas,¹³ it is generally accepted that PCs are larger than other neoplasms.^{12,14-16} Size comparisons in the literature primarily focus on PCs versus APTs, and few studies include PAs in such analyses. One example is the study by Chen et al., which evaluated characteristics of PAs, APTs, and PCs together. They reported significant differences in size among the groups, with APTs occupying an intermediate position between PAs and PCs. In the present study, the median diameter was 17 mm (13-21) for PAs and 20 mm (15-27) for APTs, showing borderline significance ($p = 0.054$). Considering that a single diameter measurement may not always accurately reflect lesion volume, we calculated lesion volumes, unlike prior studies. APTs were found to be larger in volume




Figure 2. Receiver operating characteristic curve analysis for pre-operative parathyroid hormone (area under the curve: 0.691, sensitivity: 76%. Specificity: 58%, positive predictive value: 48%, negative predictive value: 83%).




Figure 3. Receiver operating characteristic curve analysis for pre-operative calcium (area under the curve: 0.751, Sensitivity: 70%, Specificity: 69%, positive predictive value: 53%, negative predictive value: 82%).

compared to PAs (median: 996.5 mm³ vs. 734.34 mm³), but the difference was not statistically significant ($p = 0.085$). Consistent with the literature, these results suggest that APTs tend to be larger. However, no definitive cutoff values for size or volume could be

established for APTs, indicating that pre-operative differentiation of APTs from PAs based on size alone remains challenging.

PHPT often presents with mild hypercalcemia (serum Ca < 11 mg/dL).¹⁷ Studies comparing Ca and PTH levels in patients with PA, APT, and PC have shown that the increase in Ca and PTH levels in patients with APT is not as high as that in patients with PC.¹⁸⁻²⁰ Cakir et al. and McCoy et al. compared APTs with PAs and PCs and found significant differences among these three patient groups in terms of PTH and Ca levels.^{10,18} Cakir et al. grouped APT and PC under “aggressive tumors” and compared them with PA. Their analysis identified a cutoff value for Ca (12.45 mg/dL), but not for PTH levels. In the present study, significant differences in Ca and PTH levels were observed between patients with APT and those with PA, with cutoff values established as 11.50 mg/dL and 180 pg/mL, respectively. We attribute the higher Ca cutoff reported by Cakir et al. to their inclusion of patients with PC, rendering their cutoff of 12.45 mg/dL less applicable for patients with APTs. To the best of our knowledge, the present study is the first to report cutoff values for distinguishing APTs from PAs.

Data on the most common gland from which PAs originate, and therefore, the most common localization of PAs, are very limited. In the present study, a significant difference was observed between the anatomical localization of APTs and PAs. APTs were predominantly located in the left superior and right inferior poles, whereas PAs showed a homogeneous distribution.

The coexistence of thyroid disease with PHPT has been reported in 22-70% of cases since the 1950s.²¹⁻²³ In the present study, concurrent thyroid surgery (lobectomy or total thyroidectomy) was performed in 58% of APT patients versus 32% of PA patients. The difference between these rates was statistically significant. This finding is consistent with the result reported by Chen et al.¹² We hypothesize that the higher rate of thyroid surgery in APT patients may be due to concomitant thyroid pathology as well as tumor adherence to thyroid tissue. Therefore, in patients with a strong suspicion of APT, the potential need for concurrent thyroidectomy should be considered pre-operatively, and patients should be informed accordingly.

In the present study, patients were evaluated postoperatively for transient/permanent nerve injury, transient/permanent hypoparathyroidism, and HBS. No studies have compared nerve injury rates between APT and PA patients thus far. However, Gurrado et al. reported transient nerve injury in 3.16% and permanent nerve

injury in 1.4% of APT patients.¹⁵ Tang et al. documented a nerve injury rate of 14% in APT patients.²⁴ In the present study, transient nerve injury occurred in 16% of APT patients, with no permanent nerve injury observed. In PA patients, these rates were 6% and 1%, respectively. Transient nerve injury was significantly more common in APT patients compared to that in PA patients. We hypothesize that this higher risk in APT patients may stem from tumor adherence to nerves/adjacent tissues or the higher frequency of concurrent thyroid surgery. Based on these findings, we suggest that intraoperative nerve monitoring and heightened surgical caution should be considered for patients with pre-operative Ca > 11.50 mg/dL and PTH > 180 pg/mL. Gurrado et al. reported transient hypoparathyroidism in 22.3% and permanent hypoparathyroidism in 4.1% of APT patients.¹⁵ In the present study, these rates were 8% and 1%, respectively, with no significant difference compared to these rates in PA patients.

HBS is characterized by post-operative hypocalcemia, hypophosphatemia, and prolonged hypocalcemia.²⁵ It occurs in 12-25% of cases after parathyroidectomy.²⁶ Tang et al. observed HBS in 7% of APT patients in the post-operative period.²⁴ In the present study, these rates were 16% in APT patients and 3% in PA patients, and the difference was statistically significant. Based on these data, we recommend that patients with a strong suspicion of APT should be monitored more closely for the risk of HBS, and pre-operative initiation of calcitriol and oral calcium supplements should be considered to mitigate this risk.

In 2022, the WHO reclassified parathyroid tumors, defining the atypical APT diagnostic criteria as follows: cellular nests in a thickened connective tissue, tumor cells in the capsule, adherence to adjacent structures without frank invasion, band-like fibrosis, trabecular growth, increased mitotic activity (> 5/10 mm²), atypical mitotic figures, coagulative necrosis, PFIB loss, and a Ki-67 labeling index > 5%. APT cases were further stratified into PFIB-positive (PFIB+) and PFIB-negative (PFIB-) groups, with PFIB+ tumors classified as very low-risk and PFIB- as low-risk for recurrence.³ In the present study, histopathological features of APT patients were reassessed against these criteria, confirming all diagnoses. Band-like fibrosis (84%) and tumor cells within the capsule (40%) were most common, while coagulative necrosis and increased mitotic activity (> 5/10 mm²) were rare (4%). Other histopathological features were detected at varying frequencies.

To date, there are no specifically established guidelines for the post-operative follow-up of patients with

APT. While some studies, despite the low risk of recurrence, recommend close follow-up of patients independent of immunohistochemical results based on their own experience,^{27,28} other researchers advocate for close followup in patients with germline gene mutations and/or complete loss of PFIB expression.^{5,8,29} Several studies did not report any recurrence during follow-up;^{11,18,24} however, one study reported a recurrence rate of 5.3%.¹³ In the present study, median follow-up time was 4.8 years. During this period, no recurrence was observed in either group. However, persistent disease necessitating a second operation was observed in three patients with PA and two patients with APT. In patients with persistent PA, the adenomas were not completely removed during the first operation. In APT patients, persistent disease was attributed to incomplete initial tumor resection or additional adenomas. Although no significant difference was detected between the two groups in terms of persistent disease and recurrence, we recommend closer follow-up for patients with atypical tumors compared to that for those with adenomas; this recommendation is due to the serious histopathological findings observed in the atypical tumor group.

Conclusions

Pre-operative identification of APTs is critical. Ca > 11.50 mg/dL and PTH > 180 pg/mL values can be used to differentiate APTs from benign adenomas. Patients with these values should be provided with detailed pre-operative counseling regarding possible concomitant thyroid surgery, nerve injury, and HBS complications. No recurrence was observed in our follow-up, we recommend stricter and more vigilant monitoring for APTs compared to adenomas due to their concerning histopathological features.

Limitation

The limitations of this study include its retrospective nature, which inevitably leads to data loss. Other limitations are that the surgical treatment and histopathological examination were not performed by the same physicians, and the cases could not be grouped according to PFIB status.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of human subjects and animals. The authors declare to have followed the ethical standards of the relevant experimentation committee, according to the World Medical Association and the Declaration of Helsinki. The procedures were approved by the institutional Ethics Committee.

Confidentiality, informed consent, and ethical approval. The authors have obtained approval from the Ethics Committee for the analysis of routinely collected and anonymized clinical data; therefore, individual informed consent was not required. Relevant ethical recommendations have been followed.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing or creation of the content of this manuscript.

References

1. Rodgers SE, Perrier ND. Parathyroid carcinoma. *Curr Opin Oncol.* 2006;18:16-22.
2. Cetani F, Marcocci C, Torregrossa L, Pardi E. Atypical parathyroid adenomas: challenging lesions in the differential diagnosis of endocrine tumors. *Endocr Relat Cancer.* 2019;26:R441-64.
3. Erickson LA, Mete O, Juhlin CC, Perren A, Gill AJ. Overview of the 2022 WHO classification of parathyroid tumors. *Endocr Pathol.* 2022;33:64-89.
4. American Joint Committee on Cancer. *AJCC Cancer Staging Form Supplement to the AJCC Cancer Staging Manual.* 8th ed. Berlin: Springer; 2018.
5. Gill AJ, Lim G, Cheung VK, Andrici J, Perry-keene JL, Paik J, et al. Parafibromin-deficient (HPT-JT type, CDC73 mutated) parathyroid tumors demonstrate distinctive morphologic features. *Am J Surg Pathol.* 2019;43:35-46.
6. Juhlin CC, Nilsson IL, Lagerstedt-Robinson K, Stenman A, Bränström R, Tham E, et al. Parafibromin immunostainings of parathyroid tumors in clinical routine: a near-decade experience from a tertiary center. *Mod Pathol.* 2019;32:1082-94.
7. Juhlin CC, Nilsson IL, Johansson K, Haglund F, Villablanca A, Hoog A, et al. Parafibromin and APC as screening markers for malignant potential in atypical parathyroid adenomas. *Endocr Pathol.* 2010;21:166-77.
8. Kruijff S, Sidhu SB, Sywak MS, Gill AJ, Delbridge LW. Negative parafibromin staining predicts malignant behavior in atypical parathyroid adenomas. *Ann Surg Oncol.* 2014;21:426-33.
9. Saponaro F, Pardi E, Mazoni L, Borsari S, Torregrossa L, Apicella M, et al. Do patients with atypical parathyroid adenoma need close follow-up? *J Clin Endocrinol Metab.* 2021;106:4565-79.
10. Cakir B, Polat SB, Kilic M, Ozdemir D, Aydin C, Süngü N, et al. Evaluation of preoperative ultrasonographic and biochemical features of patients with aggressive parathyroid disease: is there a reliable predictive marker? *Arch Endocrinol Metab.* 2016;60:537-44.
11. Galani A, Morandi R, Dimko M, Molfino S, Baronchelli C, Lai S, et al. Atypical parathyroid adenoma: clinical and anatomical pathologic features. *World J Surg Oncol.* 2021;19:19.
12. Chen Y, Song A, Nie M, Jiang Y, Li M, Xia W, et al. Clinical and genetic analysis of atypical parathyroid adenoma compared with parathyroid carcinoma and benign lesions in a Chinese cohort. *Front Endocrinol (Lausanne).* 2023;14:1027598.
13. Marini F, Marcucci G, Giusti F, Arvat E, Benvenga S, Bondanelli M, et al. Parathyroid carcinoma and atypical parathyroid tumor: analysis of an Italian database. *Eur J Endocrinol.* 2024;191:416-25.

14. Bae JH, Choi HJ, Lee Y, Moon MK, Park YJ, Shin CS, et al. Preoperative predictive factors for parathyroid carcinoma in patients with primary hyperparathyroidism. *J Korean Med Sci.* 2012;27:890-5.
15. Gurrado A, Pasculli A, Avenia N, Bellantone R, Boniardi M, Merante Boschini I, et al. Parathyroid retrospective analysis of neoplasms incidence (pTRANI Study): an Italian multicenter study on parathyroid carcinoma and atypical parathyroid tumour. *J Clin Med.* 2023;12:6297.
16. O'Neal P, Mowschenson P, Connolly J, Hasselgren PO. Large parathyroid tumors have an increased risk of atypia and carcinoma. *Am J Surg.* 2011;202:146-50.
17. Carroll MF, Schade DS. A practical approach to hypercalcemia. *Am Fam Physician.* 2003;67:1959-66.
18. McCoy KL, Seethala RR, Armstrong MJ, Nikiforova MN, Stang MT, Carty SE, et al. The clinical importance of parathyroid atypia: is long-term surveillance necessary? *Surgery.* 2015;158:929-35.
19. Quinn CE, Healy J, Lebastchi AH, Brown TC, Stein JE, Prasad ML, et al. Modern experience with aggressive parathyroid tumors in a high-volume New England referral center. *J Am Coll Surg.* 2015;220:1054-62.
20. Fernandez-Ranvier GG, Khanafshar E, Jensen K, Zarnegar R, Lee J, Kebebew E, et al. Parathyroid carcinoma, atypical parathyroid adenoma, or parathyromatosis? *Cancer.* 2007;110:255-64.
21. Regal M, Páramo C, Luna Cano R, Pérez Méndez LF, Sierra JM, Rodríguez I, et al. Coexistence of primary hyperparathyroidism and thyroid disease. *J Endocrinol Invest.* 1999;22:191-7.
22. Sidhu S, Campbell P. Thyroid pathology associated with primary hyperparathyroidism. *Aust N Z J Surg.* 2000;70:285-7.
23. Dell'Erba L, Baldari S, Borsato N, Bruno G, Calò-Gabrieli G, Carletto M, et al. Retrospective analysis of the association of nodular goiter with primary and secondary hyperparathyroidism. *Eur J Endocrinol.* 2001;145:429-34.
24. Tang AL, Aunins B, Chang K, Wang JC, Hagen M, Jiang L. A multi-institutional study evaluating and describing atypical parathyroid tumors discovered after parathyroidectomy. *Laryngoscope Invest Otolaryngol.* 2022;7:901-5.
25. Khan MI, Waguespack SG, Hu MI. Medical management of postsurgical hypoparathyroidism. *Endocr Pract.* 2011;17 Suppl 1:18-25.
26. Carsote M, Nistor C. Forestalling hungry bone syndrome after parathyroidectomy in patients with primary and renal hyperparathyroidism. *Diagnostics (Basel).* 2023;13:1953.
27. Kumari N, Chaudhary N, Pradhan R, Agarwal A, Krishnani N. Role of histological criteria and immunohistochemical markers in predicting risk of malignancy in parathyroid neoplasms. *Endocr Pathol.* 2016;27:87-96.
28. Hu Y, Cui M, Xia Y, Su Z, Zhang X, Liao Q, et al. The clinical features of cystic parathyroid adenoma in Chinese population: a single-center experience. *Int J Endocrinol.* 2018;2018:3745239.
29. Agarwal A, Pradhan R, Kumari N, Krishnani N, Shukla P, Gupta SK, et al. Molecular characteristics of large parathyroid adenomas. *World J Surg.* 2016;40:607-14.