

Indocyanine green clearance test as a predictor of chemotherapy liver toxicity and post-operative complications in patients with colorectal liver metastases

La prueba de aclaramiento del verde de indocianina como predictor de daño hepático inducido por quimioterapia y de complicaciones posoperatorias en pacientes con metástasis hepáticas de origen colorrectal

Sergio Cortese¹, María I. Peligros², Emma Sola², Pilar García-Alfonso³ and José M. Tellado^{1*}

¹Department of General and Digestive Surgery; ²Department of Pathology; ³Department of General Oncology. Hospital General Universitario Gregorio Marañón, Madrid, Spain

Abstract

Objective: This study aims to investigate the relationship between the pre-operative indocyanine green (ICG) test, the chemotherapy-associated liver injury (CALI), and the development of severe post-operative complications (POC) in patients operated of colorectal liver metastases (CRLMs). **Materials and methods:** Sixty-nine patients previously treated with chemotherapy and submitted to liver resection for CRLM were retrospectively studied. Two pathologists independently reviewed the pathological specimens and assessed the presence of CALI. The correlation between ICG clearance and specific pathological features was analyzed. In addition, a logistic regression analysis was performed to seek for pre-operative factors associated with severe complications. **Results:** After a mean of 10.6 ($\pm$ 7.5) chemotherapy cycles, 44 patients (63.8%) developed CALI. ICG retention rate at 15 min (ICG-R15) was not statistically different between patients with and without CALI and it could only discriminate the presence of centrilobular fibrosis. Rate of severe complications was almost 6-fold in patients with CALI compared to patients without CALI ($p = 0.024$). ICG-R15 $\geq 10\%$ was the only independent risk factor associated with severe POC at multivariable logistic regression (OR = 4.075 95%CI: 1.077-15.422, $p = 0.039$). **Conclusions:** Pre-operative ICG clearance test, although not useful to identify patients with hepatic drug toxicity, is a strong predictor for the development of severe post-hepatectomy complications.

Keywords: Drug toxicity. Liver neoplasm. Liver function test. Post-operative complications.

Resumen

Objetivo: Investigar la relación entre el test de aclaramiento del verde de indocianina (ICG) preoperatorio, las alteraciones patológicas derivadas de la quimioterapia (CALI) y el desarrollo de complicaciones posoperatorias en los pacientes sometidos a resección hepática por metástasis de cáncer colorrectal (MCCR). **Material y métodos:** Sesenta y nueve pacientes previamente tratados con quimioterapia y operados de MCCR se estudiaron de manera retrospectiva. Dos patólogos revisaron independientemente el parénquima hepático no tumoral de los especímenes y determinaron la presencia de daño quimio-inducido. Se analizó la correlación entre el aclaramiento de ICG y las diferentes alteraciones anatómo-patológicas encontradas. Además, se realizó un análisis de regresión logística para identificar los factores preoperatorios asociados con las complicaciones posoperatorias.

*Correspondence:

Jose M. Tellado

E-mail: josemaria.tellado@salud.madrid.org

Date of reception: 08-04-2021

Date of acceptance: 25-05-2021

DOI: 10.24875/CIRUE.M21000482

Cir Cir (Eng). 2022;90(3):306-314

Contents available at PubMed

www.cirugiaycirujanos.com

2444-0507/© 2021 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Resultados: Tras una media de 10.6 ($\pm$ 7.5) ciclos de quimioterapia, 44 pacientes (63.8%) desarrollaron CALI. La tasa de retención de ICG a los 15 minutos (ICG-R15) no fue estadísticamente diferente entre los pacientes con y sin CALI y solo pudo discriminar la presencia de fibrosis centrolobulillar. La tasa de complicaciones severas posoperatorias fue 6 veces superior en los pacientes con CALI, comparada con aquella de los pacientes sin CALI ($p = 0.024$). Un ICG-R15 $\geq 10\%$ fue el único factor de riesgo independiente asociado a complicaciones severas (OR = 4.075 95%CI: 1.077-15.422, $p = 0.039$). **Conclusiones:** La prueba preoperatoria de aclaramiento del ICG, a pesar de no identificar eficazmente los pacientes con daño por quimioterapia, es un fuerte predictor de desarrollo de complicaciones severas posoperatorias.

Palabras clave: Toxicidad por fármacos. Metástasis hepáticas. Test de función hepática. Complicaciones posoperatorias.

Introduction

Surgical resection is the only treatment option that can offer long-lasting survival in patients with colorectal liver metastases (CRLMs)¹. Modern chemotherapeutic agents have led to an increase of resectability in those patients with initially unresectable CRLM². However, the advantages of the neoadjuvant treatment may be hampered by the side effects of chemotherapy on the non-tumoral liver. Different macroscopic and histopathological entities related to the use of these drugs have been documented. Oxaliplatin-based regimens are associated with hepatic sinusoidal congestion ("blue liver"), caused by the rupture of the sinusoidal membrane and the collagenization of perisinusoidal space³, while various degrees of liver steatosis and steatohepatitis ("yellow liver") may follow irinotecan-based schemes^{4,5}.

Sinusoidal obstructive syndrome (SOS) and chemotherapy-associated steatohepatitis (CASH) have been associated with a higher post-operative mortality, morbidity, and in-hospital stay after major hepatectomy^{6,7}.

An adequate pre-operative evaluation of the liver functional reserve and, hypothetically, of parenchymal injury, should be the cornerstone of prevention from the risk of post-operative liver-related complications. Indocyanine green (ICG) retention rate at 15 min (ICG-R15) is a validated test of hepatic function, useful to calculate the security volume threshold before hepatic resection in patients with underlying cirrhosis^{8,9} and in other different settings¹⁰.

The primary goal of this study is to study the relationship between pre-operative ICG-R15 and the severity of chemotherapy-associated liver injury (CALI). The secondary objective is to investigate the predictive value of pre-operative variables on the development of post-operative complications (POCs).

Materials and methods

Patient's selection

For the purposes of this study, we selected 69 patients out of our prospectively maintained database of consecutive patients who underwent curative resection for CRLM at our hospital over a 7-year period, on the basis of the following selection criteria: (1) availability of sufficient non-tumoral liver parenchyma for pathological analysis; (2) preoperatively recorded ICG test values; (3) no chronic underlying liver disease, and (4) known data about pre-operative chemotherapy (adjuvant post-colectomy and/or neoadjuvant before hepatic surgery). We discarded patients submitted to synchronic liver and bowel surgery (who could show morbidity not related to the liver procedure) and with previous portal embolization (who could present parenchymal and vascular alterations not related to chemotherapy). The study was conducted according to the principles of the Declaration of Helsinki and was approved by the local ethics committee.

Pre-operative chemotherapy

Chemotherapy protocols were simplified into three groups: (1) oxaliplatin group (who received FOLFOX); (2) irinotecan group (who received FOLFIRI); and (3) sequential group (patients who switched from FOLFOX to FOLFIRI or vice versa). In about two-thirds of the patients, a monoclonal antibody (cetuximab or bevacizumab) was included in the chemotherapeutic regimen. Clinical decision-making of each case went throughout a weekly discussion at the Multidisciplinary Committee of our Hospital devoted to digestive cancer.

Pre-operative planning

Number, size, and location of lesions together with liver vascular inflow and outflow were evaluated with

a pre-operative computed tomography (CT) scan performed 4-6 weeks before surgery; a magnetic resonance was realized in all the cases of ill-defined lesions. A 3D liver reconstruction and a volumetric study based on the pre-operative CT were performed. In major hepatectomies, a future liver remnant volume of at least 30% of the total liver volume was considered adequate.

From serum aspartate aminotransferase (AST) level and platelet count recorded before surgery, we calculated the AST-to-platelet ratio index (APRI) score¹¹, to be correlated with pathologic findings.

ICG test was performed the day before surgery, injecting intravenously a bolus of 0.5 mg/kg of body weight of the fluorescent dye ICG (ICG-PULSION, Germany) and recording the retention rate at 15 min (ICG-R15 as a percentage) during the hepatic clearance time by means of pulse spectrophotometry (PULSION Medical System, Germany). An impaired ICG clearance did not preclude the performance of major resections (≥ 3 Couinaud segments). However, whenever feasible, parenchymal sparing resections were preferred, trying to assure an oncological tumor-free margin of at least 1 mm¹².

Surgical procedure

In all patients, we performed a thoroughly intraoperative restaging ultrasound (US) (MyLab™70 XVG, Esaote Platform, Italy). A contrast-enhanced US (SonoVue, Bracco, Italy) was realized in the cases of doubtful or isoechoic lesions.

Liver transection was generally realized with the complementary use of Cavitron US surgical aspirator (CUSA, Tyco Healthcare, USA) and LigaSure (Covidien, UK). Low central venous pressure (< 4 mmHg) was maintained during the transection phase to minimize venous bleeding; intermittent pedicle clamping (Pringle maneuver) was used on demand. Low-intensity radio-frequency (TissueLink, Medical Inc., USA) was used to cauterize the parenchymal transection surface.

Pathological analysis

Histopathological injury in the non-tumoral liver parenchyma was assessed reviewing archival pathological specimens (previously formalin-fixed, paraffin-embedded, and stained with hematoxylin/eosin and Masson's trichrome), selecting non-neoplastic areas distant from the tumor. All the specimens were




Figure 1. **A:** low-power examination of the liver reveals sinusoidal dilatation of centrilobular and mediolobular areas. **B:** at $\times 4$ is shown nodular hyperplasia areas delimited by portal tracts and atrophic hepatocytes. **C:** high-power examination reveals fibrous tissue (arrow) around centrilobular vein. **D:** low-power photomicrograph shows severe macrovesicular steatosis. (**A**, **B**, and **D** on hematoxylin and eosin stain; **C** on Trichrome Masson stain).

reviewed independently by two pathologists (I.P. and E.S.) who were unaware of any clinical data related to pre-operative chemotherapy.

SOS (Fig. 1A) was punctuated using the pathological score reported by Rubbia-Brandt et al.³ and considered pathological when it reached a Grade 2 or 3 over 3. The presence of nodular regenerative hyperplasia (NRH) (Fig. 1B) and centrilobular vein fibrosis (CVF) (Fig. 1C) was also assessed¹³. CASH (Fig. 1D) was evaluated by means of the non-alcoholic fatty liver disease (NAFLD) activity score, as reported by Kleiner et al.¹⁴, and classified as pathological with a score of 4 or superior. To simplify the analysis, CALI was considered to be present if the non-tumoral parenchyma showed at least a pathological SOS, a pathological NAFLD, or the association of CVF and NRH.

Post-operative evaluation

Post-operative morbidity was classified using the Clavien-Dindo scale¹⁵, and major complications were classified as Clavien-Dindo III-V. An abdominal contrast-enhanced CT was realized in every case of suspected surgical complication. Post-hepatectomy liver failure (PHLF) was defined using the "50-50" criteria (prothrombin time $< 50\%$ and total bilirubin 50 micromol/L at post-operative day 5)¹⁶. Bile leakage was defined as bile-stained liquid in the abdominal drainage at any moment after hepatectomy,

after percutaneous puncture, or found during relaparotomy. Post-operative mortality was considered in the 90 days following surgery.

Statistical analysis

The statistical analysis was performed using SPSS software (version 25.0, IBM, USA). To compare variables between groups, Student's t, two-tailed Mann-Whitney, Chi-square, and Fisher's exact tests were used when appropriate. All predictors with $p < 0.10$ by univariate analysis were considered in the multivariate model. A multivariate logistic regression analysis was performed to identify independent factors. Prediction accuracy was evaluated with the area under the receiver operating characteristic curve. $p < 0.05$ was considered to indicate a statistically significant difference.

Results

Patient's characteristics

The clinicopathological characteristic and surgical procedures of the entire cohort are listed in table 1. Of the 69 patients analyzed, 58.6% were male and 41.4% were female. Mean age was 59.6 ($\pm$ 12.1) years. Liver metastasis was synchronous with the primary tumor in 63.8% of cases, and patients presented an average of 4.9 ($\pm$ 5.5) liver nodules. Fifty-nine patients (85.5%) received FOLFOX, 2 patients (2.9%) received FOLFIRI, and 8 patients (11.6%) received both regimens sequentially. A median of nine chemotherapy cycles (interquartile range 5.0-14.0) were delivered in each patient with a median interval before surgery of 6.4 weeks (interquartile range 5.0-11.6 weeks). Major resections represented 47.8% of the total procedures, minor resections (1 or 2 Couinaud segments) 37.7%, and non-anatomical wedge resections represented 14.5%. Radiofrequency ablation (RFA) was associated to liver resection in 19 cases (27.5%).

Pathological analysis and prediction of CALI

Pathological analysis of non-tumoral liver detected CALI in 44 liver specimens (63.8%), while the remaining 25 (36.2%) showed a normal parenchyma or did not fulfill the criteria of CALI. The subtypes of pathological injuries described are shown in figure 2.

Table 1. Patient's characteristics

Characteristic	Value
Total no. of patients	69
Sex (male/female)	40/29
Age, years	59.6 (12.1)
Synchronous metastases (%)	44 (63.8)
Extrahepatic disease (%)	14 (20.3)
No. of liver metastases	4.9 (5.5)
Pre-operative CEA $\geq$ 10 ng/ml (%)	20 (29.0)
Maximum size of metastases, mm	32.2 (20.0)
Chemotherapeutic agents	
Oxaliplatin group (%)	59 (85.5)
Irinotecan group (%)	2 (2.9)
Sequential group (oxaliplatin and irinotecan) (%)	8 (11.6)
Use of bevacizumab (%)	35 (50.7)
Use of cetuximab (%)	15 (21.7)
No. of chemotherapy cycles, median (IQR)	9.0 (5.0-14.0)
Chemotherapy washout period, weeks, median (IQR)	6.4 (5.0-11.6)
Type of hepatic resection	
Major resection ($\geq$ 3 Couinaud segments) (%)	33 (47.8)
Minor resection (1 or 2 Couinaud segments) (%)	26 (37.7)
Wedge resection/s (%)	10 (14.5)
Associated RFA (%)	19 (27.5)

Data are expressed as mean (standard deviation), median (interquartile range, IQR), or number (percentage), when indicated.

CEA: carcinoembryonic antigen; RFA: radiofrequency ablation.




Figure 2. Proportion of patients with and without CALI. In the histogram, individual pathological features and their frequencies are specified. CALI: chemotherapy-induced liver injury; CASH: chemotherapy-associated steatohepatitis; CVF: centrilobular vein fibrosis; NRH: nodular regenerative hyperplasia; SOS: sinusoidal obstructive syndrome.

Patients with CALI were older and of male sex in a higher proportion compared to patients without CALI, although not significantly ($p = 0.071$ and $p = 0.076$, respectively) (Table 2). The presence of dyslipidemia was strongly associated with the development of CALI ($p = 0.033$), which, in turn, showed a dose-dependent

Table 2. Comparison of pre-operative characteristic between patients with and without chemotherapy-associated liver injury (CALI)

Pre-operative characteristics	Non-CALI (n = 25)	CALI (n = 44)	p
Age, years	56.2 (11.2)	61.6 (12.2)	0.071
Age ≥ 65 years (%)	8 (32.0)	21 (47.7)	0.203
Male sex (%)	11 (44.0)	29 (65.9)	0.076
Diabetes (%)	2 (8.0)	2 (4.5)	0.460
Dyslipidemia (%)	2 (8.0)	13 (29.5)	0.033
BMI ≥ 28 kg/m ² (%)	3 (12.0)	5 (11.4)	0.611
Alkaline phosphatase, U/L	104.6 (37.4)	98.4 (40.5)	0.543
Alanine aminotransferase, U/L	27.1 (11.4)	30.0 (27.4)	0.619
Aspartate aminotransferase, U/L	30.3 (14.7)	30.0 (21.7)	0.950
ICG-PDR, %/min	19.3 (6.2)	17.7 (4.2)	0.203
ICG-R15, %	7.5 (4.8)	8.9 (5.0)	0.267
ICG-R15 ≥ 10% (%)	7 (28.0)	16 (36.4)	0.479
Platelet count (10 ³ /mm ³)	198.1 (48.4)	209.3 (75.3)	0.505
APRI score	0.52 (.27)	0.51 (.33)	0.879
No. of chemotherapy cycles	9.6 (8.2)	11.2 (7.2)	0.426
≥ 0.426 chemotherapy cycles (%)	10 (40.0)	27 (61.4)	0.087
≥ 12 chemotherapy cycles (%)	6 (24.0)	21 (47.7)	0.049
Chemotherapy washout period, weeks	10.7 (14.3)	15.6 (19.7)	0.292
Use of both oxaliplatin and irinotecan (%)	4 (16.0)	4 (9.1)	0.312
Use of irinotecan (%)	4 (16.0)	6 (13.6)	0.525
Use of bevacizumab (%)	11 (44.0)	24 (54.5)	0.400
Use of cetuximab (%)	7 (28.0)	8 (18.2)	0.342

Data are expressed as mean (standard deviation), or number (percentage), when indicated. p values refer to Student's t or two-tailed Mann-Whitney for continuous variables and Chi-square or Fisher's exact tests for categorical variables.

CALI: chemotherapy-associated liver injury; BMI: body mass index; ICG-PDR: indocyanine green plasma disappearance rate; APRI: aspartate aminotransferase-to-platelet ratio index; ICG-R15: indocyanine green retention rate at 15 min.

onset, being significantly more frequent after 12 cycles of chemotherapy ($p = 0.049$).

None of the pre-operative routine laboratory tests (aminotransferases, alkaline phosphatase, and platelet count) reflected the presence of CALI. APRI score was similar between patients with and without CALI ($p = 0.879$). ICG-R15 showed a trend to be more pathological in patients with CALI ($8.9 \pm 5.0\%$), compared to patients without CALI ($7.5 \pm 4.8\%$) but without statistical significance ($p = 0.267$).

Considering separately each pathological injury described, a significant relationship could be found between ICG-R15 and the presence of CVF, but only when the pathological cutoff was set at 16% (Table 3).

CALI, ICG-R15, and post-operative outcomes

Patients with CALI experienced a higher incidence of severe POC (25.0%), compared to the group without CALI (4.0%, $p = 0.024$), being the two groups comparable with respect to tumoral burden, proportion of major resections, association with RFA, and use of laparoscopic technique (Table 4). Furthermore, in-hospital stay was significantly longer in patients with CALI (a mean of 11.0 ± 10.1 days vs. 7.3 ± 4.4 days in patients without liver injury, $p = 0.039$).

Mortality, bile leakage, red blood cell (RBC) transfusion rate, and incidence of PHLF were not different between patients with and without CALI. Patients with a pathological ICG-R15 showed higher incidence of severe POC ($p = 0.043$) and a higher rate of perioperative RBC transfusion ($p = 0.032$) compared to those with a normal ICG-R15 (Table 4). In a sub-analysis of the 33 major liver resections realized, PHLF developed in four patients among 13 with ICG-R15 $\geq 10\%$ (30.8%) and only in two out of the 20 with ICG-R15 $< 10\%$ (10.0%, $p = 0.147$). Meanwhile, no significant differences in the liver tumor status and operative procedure were seen between these two groups.

Prediction of major POC

Among all the pre-operative and surgical variables analyzed, only a pathological ICG-R15 ($\geq 10\%$) was found to be an independent predictor of severe POC, including death (OR = 4.075, 95% C.I.: 1.077-15.422, $p = 0.039$, AUC 0.738) (Table 5). Use of RFA (vs. non-use) seemed to be a protective factor against POC ($p = 0.094$ at univariate analysis) and was included in the multivariate model, but did not show significance as an independent factor ($p = 0.106$).

Discussion

CALI may play an important role in the morbidity and mortality after hepatic resection for CRLM. Since the first description of SOS by Rubbia-Brandt et al.³,

Table 3. Comparison of pathological features of chemotherapy liver toxicity and patients with two cutoffs (at 10% and at 16%) of indocyanine green retention rate (ICG-R15)

Pathological injury	ICG-R15 < 10% (n = 46)	ICG-R15 ≥ 10% (n = 23)	p	ICG-R15 < 16% (n = 62)	ICG-R15 ≥ 16% (n = 7)	p
Sinusoidal obstructive syndrome (Grade 2 o 3) (%)	23 (50.0)	11 (47.8)	0.865	31 (50.0)	3 (42.9)	0.517
Centrilobular vein fibrosis (%)	4 (8.7)	5 (21.7)	0.129	6 (9.7)	3 (42.9)	0.042
Nodular regenerative hyperplasia (%)	7 (15.2)	5 (21.7)	0.500	9 (14.5)	3 (42.9)	0.095
CASH (NAFLD activity score ≥ 4) (%)	8 (17.4)	6 (26.1)	0.397	12 (19.4)	2 (28.6)	0.436

Data are expressed as number (percentage). p values refer to Chi-square or Fisher's exact tests.

CASH: chemotherapy-associated steatohepatitis; NAFLD: non-alcoholic fatty liver disease; ICG-R15: indocyanine green retention rate at 15 min.

Table 4. Comparison of intraoperative and post-operative outcomes in patients with and without CALI and between those with a normal versus ICG-R15

Characteristic	Non-CALI (n = 25)	CALI (n = 44)	p	ICG R15 < 10% (n = 46)	ICG R15 ≥ 10% (n = 23)	p
Tumor maximum diameter, mm	28.4 (16.5)	34.3 (21.6)	0.241	31.8 (21.7)	33.0 (16.5)	0.817
No. of metastases	6.2 (7.1)	4.1 (4.2)	0.116	4.7 (5.7)	5.2 (5.1)	0.702
Major resections (≥ 3 Couinaud segments) (%)	11 (44.0)	22 (50.0)	0.632	20 (43.5)	13 (56.5)	0.307
Laparoscopy (%)	1 (4.0)	6 (13.6)	0.199	5 (10.9)	2 (8.7)	0.571
Pringle maneuver, minutes	18.7 (9.4)	17.9 (11.7)	0.879	17.3 (10.1)	21.5 (11.3)	0.490
Associated RFA (%)	9 (36.0)	10 (22.7)	0.235	12 (26.1)	7 (30.4)	0.703
Minor complications (Clavien-Dindo I-II) (%)	5 (20.0)	6 (13.6)	0.488	6 (13.0)	5 (21.7)	0.352
Severe complications (Clavien-Dindo III-V) (%)	1 (4.0)	11 (25.0)	0.024	5 (10.9)	7 (30.4)	0.043
Bile leakage (%)	2 (8.0)	4 (9.1)	0.625	4 (8.7)	2 (8.7)	0.686
Post-operative mortality (90 days) (%)	0 (0)	1 (2.3)	0.453	0 (0)	1 (4.3)	0.333
Liver failure (%)						
All resections (n = 69)	2 (8.0)	5 (11.4)	0.501	3 (6.5)	4 (14.4)	0.161
Major resections (n = 33)	2 (18.2)	4 (18.2)	0.671	2 (10.0)	4 (30.8)	0.147
RBC transfusion (%)	2 (8.0)	7 (13.6)	0.294	3 (6.5)	6 (26.1)	0.032
In-hospital stay, days	7.3 (4.4)	11 (10.1)	0.039	8.9 (9.1)	11.3 (7.2)	0.279

Characteristics related to tumoral burden and surgical technique are also compared. p values refer to Student's t or two-tailed Mann-Whitney for continuous variables and Chi-square or Fisher's exact tests for categorical variables. Data are expressed as mean (standard deviation) or number (percentage), when indicated.

CALI: chemotherapy-associated liver injury; ICG-R15: pathological indocyanine green retention rate; RFA: radiofrequency ablation; RBC: red blood cell.

additional features of morphological change in non-tumoral liver due to oxaliplatin have been documented, such as centrilobular and perisinusoidal fibrosis, peliosis, and nodular hyperplasia¹³. Subsequently, other authors pointed out the usefulness of semi-quantitative grading of anti-CD34 antibodies (a marker of sinusoidal capillarization)¹⁷, and of nuclear proliferation markers¹⁸ as measurable hallmarks of CALI.

Irinotecan-induced steatohepatitis has been associated with higher post-operative mortality⁷, while the presence of SOS can be the cause of higher morbidity¹⁹, transfusion rate²⁰, and liver dysfunction (ascites and liver failure)²¹. Despite other studies could not show a clear association between CALI and a worse post-operative outcome^{22,23}, a recent review and meta-analysis that include eight retrospective Eastern and Western publications, gathering more than 700 patients²⁴, strongly

Table 5. Univariate and multivariate analysis of the influence of pre-operative factors and surgical technique on the development of severe post-operative complications and mortality (Dindo-Clavien Grades III-V) in the study cohort (n = 69)

Variables	Univariate analysis (p)	Odds ratio	95% CI	p
Age ≥ 65 years	0.161	-	-	-
Male sex	0.368	-	-	-
Dyslipidemia	0.715	-	-	-
Diabetes	0.137	-	-	-
BMI ≥ 28 kg/m ²	0.137	-	-	-
Synchronic metastases (vs. metachronic)	0.294	-	-	-
Four or more nodules	0.803	-	-	-
More than 12 chemotherapy cycles	0.121	-	-	-
Tumor major diameter ≥ 30 mm	0.178	-	-	-
APRI score > 0.5	0.200	-	-	-
ICG-R15 ≥ 10%	0.043	4.075	1.077-15.422	0.039
Major hepatectomy	0.131	-	-	-
Non-use versus use of RFA	0.094	0.167	0.019-1.463	0.106
Open surgery (versus laparoscopy)	0.649	-	-	-

APRI: aspartate aminotransferase-to-platelet ratio index; BMI: body mass index; RFA: radiofrequency ablation.

suggest that sinusoidal dilatation and steatohepatitis are associated with severe post-operative morbidity and liver dysfunction, respectively.

In this context, it seems to be very useful to have a non-invasive diagnostic tool to predict the grade of CALI to better select and prepare especially those patients requiring a major hepatectomy. ICG clearance determination is a trustful bedside test largely employed, especially in Eastern countries, to assess hepatic functional reserve in patients with known liver disease^{8,9}. ICG is an anionic dye which, following intravenous injection, almost completely binds to plasma proteins with no extravascular distribution. Its elimination is a carrier mediated process through the biliary canaliculi and no enterohepatic circulation takes place²⁵. ICG clearance reduction in cirrhosis may be explicated by the reduction of hepatic blood flow and by a reduced uptake of the dye from the

sinusoids to the hepatocytes, as a consequence of microvascular intrahepatic changes (portovenous shunts or capillarization of sinusoidal space)²⁶. A decade ago, Krieger et al. documented the influence of chemotherapy for CRLM on ICG clearance value²⁷ but these data were not confirmed by Wakiya et al.²⁸ Recently, also, Wang et al.²⁹ pointed out the relation between pre-operative ICG-R15 value and the use of chemotherapy, although in their work, no histological study of CALI was performed on liver specimens.

In our series of 69 patients, a significant relationship could be found between ICG-R15 and the presence of CVF, which is a late-onset oxaliplatin-related injury, but only when the cutoff was set at 16%. SOS, CASH, and NRH could not be efficiently detected by ICG clearance test.

Interestingly, the presence of dyslipidemia seemed to be a predisposing factor for the development of CALI. An altered free fatty acid and cholesterol metabolism and storage, as seen in dyslipidemia, could boost the toxic effect of 5-fluorouracil (a component of both FOLFOX and FOLFIRI schemes) on hepatocyte mitochondria³⁰, and possibly contribute to a pro-inflammatory environment leading to steatohepatitis.

The present study shows, on the one hand, the high prevalence of CALI (almost 64%) in patients submitted to chemotherapy for CRLM and, on the other, confirms the notion that CALI is associated with a higher risk of post-operative severe complications and prolonged hospital stay.

Surprisingly, although ICG-R15 was only marginally correlated to the severity of CALI, it showed to be the best predictor of severe POC. Furthermore, perioperative RBC transfusion rate was significantly higher in patients with an impaired ICG-R15.

In our usual practice for CRLM, we do not consider a pathological ICG-R15 as a contraindication to perform major resections. In a sub-analysis performed considering exclusively major hepatectomies, we found a trend toward a higher incidence of liver failure among patients with a pathological ICG-R15 compared to those with a normal one, but without statistical significance.

These data should be interpreted with caution due to the limited number of events in these groups and a possible problem of underpower, but suggest that parenchymal sparing procedures and a meticulous technical execution should be advisable in patients with a pre-operative ICG-R15 ≥ 10%.

In a retrospective series of 161 patients with CRLM who received pre-operative chemotherapy, the group of Makuuchi³¹ performed 37 major hepatectomies in patients with an ICG-R15 $\geq 10\%$, although theoretically assuring at least 60% of the future liver remnant volume by means of portal vein embolization. In this subgroup of patients with marginal liver functional reserve, the authors observed significantly poorer blood test values associated with liver dysfunction, higher amount of blood loss, and significantly higher total morbidity.

We acknowledge some limitations of our study, mainly caused by the retrospective design: first, the rather wide variability between the last chemotherapy cycle and surgery. This could be explained by the fact that more than one-third of our patients presented liver metachronic disease and, for many of them, the last chemotherapy cycle dated back to the end of adjuvant treatment after colonic surgery. It has been documented that, after 2-4 weeks of chemotherapy cessation, ICG values improve gradually³², while histological injury may persist months afterward³³.

Another relative drawback is due to the fragmentation of pathological features of chemotherapy toxicity that corresponds to various and often unrelated histological changes, dependent on specific drugs. The synthesis of these features in a single variable (CALI) is arbitrary but, we believe, useful to interpret the impact of liver injury in a clinical setting.

Conclusions

In patients with intense exposure to chemotherapy undergoing liver resection for CRLM, pre-operative ICG clearance test, although does not directly measure the presence of CALI, is useful to predict post-operative severe complications.

Funding

The authors declare have no funding.

Conflicts of interest

The authors declare have no conflicts of interest.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. Right to privacy and informed consent. The authors have obtained approval from the Ethics Committee for analysis and publication of routinely acquired clinical data and informed consent was not required for this retrospective observational study.

References

1. Kopetz S, Chang GJ, Overman MJ, Eng C, Sargent DJ, Larson DW, et al. Improved survival in metastatic colorectal cancer is associated with adoption of hepatic resection and improved chemotherapy. *J Clin Oncol.* 2009;27:3677-83.
2. Nordlinger B, Sorbye H, Glimelius B, Poston GJ, Schlag PM, Rougier P, et al. Perioperative chemotherapy with FOLFOX4 and surgery versus surgery alone for resectable liver metastases from colorectal cancer (EORTC Intergroup trial 40983): a randomised controlled trial. *Lancet.* 2008;371:1007-16.
3. Rubbia-Brandt L, Audard V, Sartoretto P, Roth AD, Brezault C, Le Charpentier M, et al. Severe hepatic sinusoidal obstruction associated with oxaliplatin-based chemotherapy in patients with metastatic colorectal cancer. *Ann Oncol.* 2004;15:460-6.
4. Pawlik TM, Olino K, Gleisner AL, Torbenson M, Schulick R, Choti MA. Preoperative chemotherapy for colorectal liver metastases: impact on hepatic histology and postoperative outcome. *J Gastrointest Surg.* 2007;11:860-8.
5. Robinson SM, Wilson CH, Burt AD, Manas DM, White SA. Chemotherapy-associated liver injury in patients with colorectal liver metastases: a systematic review and meta-analysis. *Ann Surg Oncol.* 2012;19:4287-99.
6. Nakano H, Oussoultzoglou E, Rosso E, Casnedi S, Chenard-Neu MP, Dufour P, et al. Sinusoidal injury increases morbidity after major hepatectomy in patients with colorectal liver metastases receiving preoperative chemotherapy. *Ann Surg.* 2008;247:118-24.
7. Vauthey JN, Pawlik TM, Ribero D, Wu TT, Zorzi D, Hoff PM, et al. Chemotherapy regimen predicts steatohepatitis and an increase in 90-day mortality after surgery for hepatic colorectal metastases. *J Clin Oncol.* 2006;24:2065-72.
8. Miyagawa S, Makuuchi M, Kawasaki S, Kakazu T. Criteria for safe hepatic resection. *Am J Surg.* 1995;169:589-94.
9. Kubota K, Makuuchi M, Kusaka K, Kobayashi T, Miki K, Hasegawa K, et al. Measurement of liver volume and hepatic functional reserve as a guide to decision-making in resectional surgery for hepatic tumors. *Hepatology.* 1997;26:1176-81.
10. Imamura H, Seyama Y, Kokudo N, Maema A, Sugawara Y, Sano K, et al. One thousand fifty-six hepatectomies without mortality in 8 years. *Arch Surg.* 2003;138:1198-206.
11. Wai CT, Greenon JK, Fontana RJ, Kalbfleisch JD, Marrero JA, Conjeevaram HS, et al. A simple noninvasive index can predict both significant fibrosis and cirrhosis in patients with chronic hepatitis C. *Hepatology.* 2003;38:518-26.
12. Vandeweyer D, Neo EL, Chen JW, Maddern GJ, Wilson TG, Padbury RT. Influence of resection margin on survival in hepatic resections for colorectal liver metastases. *HPB (Oxford).* 2009;11:499-504.
13. Rubbia-Brandt L, Lauwers GY, Wang H, Majno PE, Tanabe K, Zhu AX, et al. Sinusoidal obstruction syndrome and nodular regenerative hyperplasia are frequent oxaliplatin-associated liver lesions and partially prevented by bevacizumab in patients with hepatic colorectal metastasis. *Histopathology.* 2010;56:430-9.
14. Kleiner DE, Brunt EM, Van Natta M, Behling C, Contos MJ, Cummings OW, et al. Design and validation of a histological scoring system for nonalcoholic fatty liver disease. *Hepatology.* 2005;41:1313-21.
15. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg.* 2004;240:205-13.
16. Balzan S, Belghiti J, Farges O, Ogata S, Sauvanet A, Delefosse D, et al. The "50-50 criteria" on postoperative day 5: an accurate predictor of liver failure and death after hepatectomy. *Ann Surg.* 2005;242:824-8.
17. Narita M, Oussoultzoglou E, Chenard MP, Fuchshuber P, Rather M, Rosso E, et al. Liver injury due to chemotherapy-induced sinusoidal obstruction syndrome is associated with sinusoidal capillarization. *Ann Surg Oncol.* 2012;19:2230-7.

18. Nalbantoglu IL, Tan BR Jr., Linehan DC, Gao F, Brunt EM. Histological features and severity of oxaliplatin-induced liver injury and clinical associations. *J Dig Dis*. 2014;15:553-60.
19. Karoui M, Penna C, Amin-Hashem M, Mitry E, Benoist S, Franc B, et al. Influence of preoperative chemotherapy on the risk of major hepatectomy for colorectal liver metastases. *Ann Surg*. 2006;243:1-7.
20. Aloia T, Sebah M, Plasse M, Karam V, Lévi F, Giacchetti S, et al. Liver histology and surgical outcomes after preoperative chemotherapy with fluorouracil plus oxaliplatin in colorectal cancer liver metastases. *J Clin Oncol*. 2006;24:4983-90.
21. Soubrane O, Brouquet A, Zalinski S, Terris B, Brézault C, Mallet V, et al. Predicting high grade lesions of sinusoidal obstruction syndrome related to oxaliplatin-based chemotherapy for colorectal liver metastases: correlation with post-hepatectomy outcome. *Ann Surg*. 2010;251:454-60.
22. Sahajpal A, Vollmer CM Jr., Dixon E, Chan EK, Wei A, Cattral MS, et al. Chemotherapy for colorectal cancer prior to liver resection for colorectal cancer hepatic metastases does not adversely affect peri-operative outcomes. *J Surg Oncol*. 2007;95:22-7.
23. Makowiec F, Möhrle S, Neeff H, Drognitz O, Illerhaus G, Opitz OG, et al. Chemotherapy, liver injury, and postoperative complications in colorectal liver metastases. *J Gastrointest Surg*. 2011;15:153-64.
24. Zhao J, van Mierlo KM, Gómez-Ramírez J, Kim H, Pilgrim CH, Pessaux P, et al. Systematic review of the influence of chemotherapy-associated liver injury on outcome after partial hepatectomy for colorectal liver metastases. *Br J Surg*. 2017;104:990-1002.
25. Schneider PD. Preoperative assessment of liver function. *Surg Clin North Am*. 2004;84:355-73.
26. Imamura H, Sano K, Sugawara Y, Kokudo N, Makuuchi M. Assessment of hepatic reserve for indication of hepatic resection: decision tree incorporating indocyanine green test. *J Hepatobiliary Pancreat Surg*. 2005;12:16-22.
27. Krieger PM, Tamandl D, Herberger B, Faybik P, Fleischmann E, Maresch J, et al. Evaluation of chemotherapy-associated liver injury in patients with colorectal cancer liver metastases using indocyanine green clearance testing. *Ann Surg Oncol*. 2011;18:1644-50.
28. Wakiya T, Kudo D, Toyoki Y, Ishido K, Kimura N, Narumi S, et al. Evaluation of the usefulness of the indocyanine green clearance test for chemotherapy-associated liver injury in patients with colorectal cancer liver metastasis. *Ann Surg Oncol*. 2014;21:167-72.
29. Wang LJ, Yan XL, Li J, Wang K, Xing BC. Indocyanine green clearance test for the preoperative assessment of chemotherapy-related hepatic injury in patients with colorectal liver metastasis. *Cancer Manag Res*. 2020;12:3237-45.
30. Zeng J, Deng S, Wang Y, Li P, Tang L, Pang Y. Specific inhibition of Acyl-CoA oxidase-1 by an acetylenic acid improves hepatic lipid and Reactive Oxygen Species (ROS) metabolism in rats fed a high fat diet. *J Biol Chem*. 2017;292:3800-9.
31. Takamoto T, Hashimoto T, Ichida A, Shimada K, Maruyama Y, Makuuchi M. Surgical strategy based on indocyanine green test for chemotherapy-associated liver injury and long-term outcome in colorectal liver metastases. *J Gastrointest Surg*. 2018;22:1077-88.
32. Takamoto T, Hashimoto T, Sano K, Maruyama Y, Inoue K, Ogata S, et al. Recovery of liver function after the cessation of preoperative chemotherapy for colorectal liver metastasis. *Ann Surg Oncol*. 2010;17:2747-55.
33. Hubert C, Fervaille C, Sempoux C, Horsmans Y, Humblet Y, Machiels JP, et al. Prevalence and clinical relevance of pathological hepatic changes occurring after neoadjuvant chemotherapy for colorectal liver metastases. *Surgery*. 2010;147:185-94.