

Evaluation of the effectiveness of human breast milk exosomes in experimental testicular torsion-detorsion injury model

Evaluación de la eficacia de los exosomas de la leche materna humana en un modelo experimental de lesión por torsión-destorsión testicular

Ünal T. Öztürk¹, Hatice S.Y. Cömert^{2*}, Gül Şalıcı², Ahmet Alver³, Sevdegül A. Mungan⁴, Neslihan Sağlam³, Şeniz Erdem³, Serdar Karakullukçu⁵, Mustafa İmamoğlu², and Haluk Sarıhan²

¹Department of Pediatric Surgery, Artvin Public Hospital, Artvin; ²Department of Pediatric Surgery; ³Department of Biochemistry; ⁴Department of Pathology; ⁵Department of Public Health. Faculty of Medicine, Karadeniz Technical University, Trabzon, Turkey

Abstract

Objective: Our aim was to examine the effectiveness of human breast milk exosomes in treating testicular torsion. **Method:** Rats were divided randomly into three equal groups. Group 1 was our control group. A torsion/detorsion model was created for the rats in Group 2. After the torsion/detorsion model, 1 mg/kg intraperitoneally human breast milk exosomes were given 30 min before detorsion in Group 3. All testicles were removed at the 6th h of the experiment, for biochemical and histopathological examination. **Results:** There was a statistically significant decrease in Group 3 MDA values compared to Group 2. When Group 3 and Group 2 SOD and CAT values were compared, it was seen that these values were higher in Group 3. Histopathologically a statistically significant difference was found in Group 3 compared to Group 1 and Group 2 in terms of ischemia/reperfusion injury. **Conclusion:** We have shown that human breast milk exosomes in testicular torsion do not completely prevent reperfusion injury but significantly preserve testicular tissue.

Keywords: Testicular torsion. Exosome. Ischemia/reperfusion.

Resumen

Objetivo: Nuestro objetivo era examinar la eficacia de los exosomas de la leche materna humana en el tratamiento de la torsión testicular. **Método:** Las ratas se estudiaron en tres grupos iguales. El grupo 1 fue nuestro grupo control. Se creó un modelo de torsión/destorsión para las ratas del grupo 2. Después del modelo de torsión/destorsión, se administró 1 mg/kg de exosomas de leche materna humana por vía intraperitoneal 30 minutos antes de la destorsión en el grupo 3. Todos los testículos se extirparon a la sexta hora del experimento para examen bioquímico e histopatológico. **Resultados:** Hubo una disminución estadísticamente significativa en los valores de MDA del grupo 3 en comparación con el grupo 2. Cuando se compararon los valores de SOD y CAT del grupo 3 y del grupo 2, se vio que estos valores eran mayores en el grupo 3. Histopatológicamente se encontró una diferencia estadísticamente significativa en el grupo 3 en comparación con el grupo 1 y el grupo 2 en términos de lesión por isquemia/reperfusión. **Conclusión:** Hemos demostrado que los exosomas de la leche materna humana en la torsión testicular no previenen completamente la lesión por reperfusión, pero preservan significativamente el tejido testicular.

Palabras clave: Torsión testicular. Exosoma. Isquemia/reperfusión.

*Correspondence:

Hatice Sonay Yalçın Cömert
E-mail: sonayyalcin@hotmail.com

Date of reception: 12-09-2023

Date of acceptance: 22-11-2023

DOI: 10.24875/CIRUE.M23000827

Cir Cir (Eng). 2025;93(2):149-155

Contents available at PubMed

www.cirugiaycirujanos.com

2444-0507/© 2023 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Testicular torsion is a surgical emergency that causes cessation of blood flow to the testis and surrounding structures due to the vertical rotation of the spermatic cord in newborns, children, and adolescents¹. The incidence in men under the age of 25 is 1/4000². Early diagnosis and surgical intervention are required to prevent infertility and subfertility³. The diagnosis of testicular torsion should be made within the first 4-6 h. Especially if there is complete spermatic cord torsion, that is if there is a 720-degree rotation, it causes irreversible ischemic damage to the testis and loss of gonads⁴.

Ischemic damage during testicular torsion occurs due to oxidative stress and inflammation due to reactive oxygen species (ROS). The primary cause of testicular damage during the ischemic injury period is associated with decreased oxygen supply⁵. Tissue reperfusion should be provided as soon as possible to preserve the viability of ischemia-induced tissues. It is known that after detorsion of the testis, reperfusion also causes cellular damage, which is associated with an increase in ROS production and neutrophil infiltration⁶. The radicals formed in this process cause peroxidation of lipids in the cell membrane, denaturation of proteins, and DNA damage⁷.

In recent years, extensive research has sought effective strategies and drugs to minimize or prevent testicular ischemia/reperfusion (I/R) injury. However, other than trying to reduce the temperature of the scrotum, this is the only method that has been successfully applied in clinical practice⁸. Therefore, there is an urgent need to develop new therapeutic agents and a standard treatment protocol to reduce mortality and morbidity rates.

Exosomes are nanoscale cell-derived extracellular vesicles secreted from various cell types and found to contain bioactive components⁹. Exosomes, which can be obtained from saliva, urine, plasma, breast milk, and many other bodily fluids, can be used for therapeutic purposes, as they provide genetic material transfer, immune regulatory roles, and intercellular communication^{6,10,11}. Exosomes are natural and, thanks to their very small size, can easily pass through anybody barrier. Considering these benefits, the therapeutic use of exosomes is inevitable¹².

In recent studies, it has been proven that there are high concentrations of exosomes in breast milk, and they have been used in various experimental studies

in the literature^{13,14}. In this study, we aimed to investigate the protective effect of human breast milk exosomes against I/R damage in testicular torsion and their usability as a treatment modality.

Method

This study was carried out with the contributions of the Departments of Biochemistry and Pathology after being planned by the Department of Pediatric Surgery of Karadeniz Technical University (KTU) Faculty of Medicine, and the ethics committee approval with protocol number 2021/10 was obtained. The surgeries were performed in the KTU Faculty of Medicine Surgical Research Laboratory. Our work was supported by the Scientific Research Projects Coordination Unit.

Study Design

Three groups of 8 Sprague Dawley rats, which were divided randomly and each weighing 250-300 gr, were used in the study. The duration of the experiment was determined as 1 day. Until the study day, the rats were housed in cages with 8 or less sawdust under standard laboratory conditions at 22 ± 2 °C constant room temperature, 60% humidity, and 12 h light/12 h dark. Rats fasted for 24 h before surgery, and only water was given. In all surgical procedures, anesthesia was administered intraperitoneally, 10 mg/kg xylazine hydrochloride (Rompun, Bayer, Turkey) and 50 mg/kg ketamine (Ketalar, Pfizer, Turkey).

Surgical Technique: The operation area was shaved and cleaned with betadine solution. An incision made from the scrotal midline was used in the surgeries. Torsion/detorsion model; the left testis was taken out and rotated 720 degrees counterclockwise. Then, the tunica albuginea was sutured with 6/0 monofilament and fixed to the scrotum so the testis would not be detorsioned again. The scrotum was closed with 4/0 silk, and torsion was performed. After 4 h, the testis was detorted and waited. Orchiectomy was performed on the left testis 2 h after the detorsion procedure.

Group 1: after anesthesia, a vertical incision was made from the scrotal midline; the left testis was taken out by cutting the gubernaculum, then placed in the scrotum in an anatomical position without torsion. The layers were closed with 4/0 silk, and the left orchiectomy was performed 6 h later.

Group 2: a torsion/detorsion model was created for the rats in this group, and left testis orchiectomy was

performed 2 h after detorsion, that is, at the 6th h of the experiment.

Group 3: a torsion/detorsion model was created for rats in this group, and 1 mg/kg ip human breast milk exosomes were given 30 min before detorsion. Left orchiectomy was performed 2 h after detorsion, at the 6th h of the experiment.

The orchiectomy material was divided into two parts. Half of the obtained testicular tissue was immediately frozen with liquid nitrogen and stored at -80 degrees until biochemical analysis. The other half was reserved for histopathological examinations.

Human breast milk collection and exosome isolation

Approximately 200-300 mL of fresh breast milk samples were collected from healthy lactating mothers who applied to the KTU University, Faculty of Medicine, Pediatric Surgery Clinic (Trabzon, Turkey). Milk samples were frozen immediately and kept at -80 °C until used. Exosomes were isolated from breast milk by differential centrifugation and ultracentrifugation. All centrifugations were performed at 4 °C, and breast milk was centrifuged at 3000×g (F0650 rotor, Beckman Coulter Instruments, USA) for 10 min to separate the milk lipid layer. The supernatant was transferred to a new tube, and centrifugation was performed again at 5,000 ×g (F0650 rotor, Beckman Coulter Instruments, USA) for 30 min to eliminate large cellular debris and remaining lipids. Then, the new supernatant was transferred to a new tube, and centrifugation was performed again at 12,000 ×g (F0650 rotor, Beckman Coulter Instruments, USA) for 10 min to remove smaller cellular debris and apoptotic cells. The exosomes were obtained from the new centrifuged milk supernatant by ultracentrifugation at 110,000 ×g for 2 h at 4°C using a JS 24.15 rotor (Beckman Coulter Instruments, USA). After removing the exosome-free supernatant, the crude exosome pellet was suspended in 1000 µL phosphate-buffered saline (PBS, pH 7.4) in a 1.5 mL Eppendorf tube and stored at -80 °C until used for the experiment.

Human breast milk-derived exosome protein extraction and confirmation by western blot

Human breast milk-derived exosomes were lysed in RIPA lysis solution (SE3924401, SERVA) containing

protease and phosphatase inhibitor cocktail (PPC1010, SIGMA) was added, and exosomes were kept on ice for 15 min. Afterward, the protein was obtained by centrifugation at 15000 xg, +4°C for 15 min, and its concentrations were analyzed using the Bicinchoninic acid (BCA) kit (LOT: UD282967, Thermo Scientific, USA) and analyzed for total CD63 (positive exosome membrane marker) and calnexin (negative marker) protein expression. Lysate was separated by SDS-PAGE and transferred onto a nitrocellulose membrane. The membranes were probed with antibodies and detected using enhanced chemiluminescence detection. Primary antibodies were as follows: anti-CD63 (1:1000; Abcam, ab68418); and anti-calnexin (1:1000; Abcam, ab213243). The secondary antibody was horseradish peroxidase (HRP)-conjugated goat anti-rabbit (1:5000; Abcam, ab97200).

The protein concentration of the breast milk exosomes used in the experiment was adjusted to 1 µg/mL by the BCA kit.

Biochemical analysis

Testicular tissue was homogenized with 2 mL of cold homogenization buffer in a test tube for about 20 s until the tissues were completely disintegrated and then centrifuged at 3,000 rpm for 10 min. 750 µL of supernatant was placed in 1.5 mL Eppendorf. 600 µL of ethanol-chloroform (2:3) mixture was added to it. It was centrifuged at 10,000 xg for 30 min at +4°C. After centrifugation, protein determination, superoxide dismutase (SOD), and catalase (CAT) activity were measured in the protein phases on the upper part of the Eppendorf. SOD enzyme activity was performed by modifying the method developed by Sun and Oberley¹⁵. The Aebi method was used to determine CAT enzyme activity¹⁶.

Testicular tissue was homogenized with 2 mL of cold homogenization buffer in a test tube for about 30 s until the tissues were completely disintegrated and then centrifuged at 3,000 rpm for 10 min. The supernatant was taken, and the Malondialdehyde (MDA) level was determined. This method was carried out by modifying the method based on measuring the absorbance at 532 nm of the color of the complex formed by MDA with thiobarbituric acid in an acidic medium¹⁷.

Histopathological assay

Tissues were fixed in Bovine solution for 72 h. Fixed tissue pieces were dehydrated by passing through

70%, 90%, 96%, and 100% graded alcohol series. It was then cleared by passing through a xylene solution. After the paraffin blocks of the tissues were prepared, 5 μ m thick sections were taken with a fully automatic microtome. Sections were stained with hematoxylin and eosin (H&E) after deparaffinization. The H&E stained slides were evaluated histopathologically with a light microscope (Olympus BX 51; Olympus Optical Co, Ltd, Tokyo, Japan), by an experienced pathologist unaware of the study groups. Damage to testicular tissues was graded by the Johnsen Testicular Biopsy Score (JTBS). In this scoring system, testicular tissues were evaluated semi-quantitatively in 5 high power fields (HPF) at x200 magnification by light microscopy.

Statistical analysis

SPSS 23.0 statistical package program was used in the analysis of the data. Descriptive statistics of the evaluation results were given as median and quartiles (Q1-Q3). The suitability of the data for normal distribution was checked with the Kolmogorov-Smirnov test. The Kruskal-Wallis test was used to compare the numerical values of three independent groups. A statistically significant level was taken as $p < 0.05$ in the Kruskal-Wallis test. Since significant results were found in the Kruskal-Wallis test, Bonferroni correction was made when making pairwise comparisons. Mann-Whitney U test was used and $p < 0.017$ ($0.05/3$) was considered significant.

Results

Confirmation of exosomes from mother milk

The presence of exosomes from human breast milk was confirmed by Western blot for the exosome membrane markers cluster of differentiation 63 (CD63) and the negative marker calnexin (Fig. 1).

Biochemical analysis

When the tissue MDA, SOD, and CAT values of the three groups were compared, a statistically significant difference was found which is shown in table 1.

When the pairwise MDA values of the groups were compared, a statistically significant difference was found between group 1-group 2 and group 2-group 3




Figure 1. Confirmation of mother-breast milk exosomes by Western Blot.

($p < 0.05$). There was no significant difference between group 1 and group 3.

When the pairwise SOD values of the groups were compared, a statistically significant difference was found between group 2 and group 3 ($p < 0.05$). There was no significant difference between group 1-group 2 and group 1-group 3.

When the pairwise CAT values of the groups were compared, a statistically significant difference was found between group 1 and group 2 ($p < 0.05$). There was no significant difference between group 1-group 3 and group 2-group 3. A pairwise comparison of tissue MDA, SOD, and CAT values is given in table 1.

Histopathological assay

When the left testis JTBS of different groups were compared histopathologically, the difference between them was found to be statistically significant. Values are given in table 2.

A statistically significant difference was found between group 1-group 2 and group 2-group 3 when comparing the JTBS values of the groups in pairs. There was no significant difference between Group 1 and group 3 ($p < 0.05$). Values are given in table 2. Histopathological images in all three groups are shown in figure 2.

Discussion

The primary pathology in testicular torsion is venous occlusion that causes testicular edema and pain. However, this venous occlusion causes hypoxia in the




Figure 2. A: sertoli cell, spermatogonia (†), and Spermatocyte (▲) in testicle of group 1. B: diffuse fresh bleeding in the interstitial area and exfoliated necrotic cells in seminiferous tubule lumen, few ghost germinal epithelial cells in the left testicular tubule lumen belonging to group 2 (▲) (H&E $\times 100$). C: Spermatids (†) and spermatozoons (▲) in the testis belonging to group 3.

Table 1. The mean values of the biochemical results of the groups

Parameters	Group 1 Median (Q1-Q3)	Group 2 Median (Q1-Q3)	Group 3 Median (Q1-Q3)	p*	Post-hoc**
MDA (nmol/mg protein)	28.45 (21.03-34.73)	68.74 (44.11-91.00)	22.38 (19.68-22.52)	< 0.001	p ¹⁻² = 0.015 p ²⁻³ < 0.001
SOD (U/mg protein)	9.83 (8.67-10.65)	10.86 (9.21-13.96)	7.30 (6.78-9.46)	0.012	p ²⁻³ = 0.004
CAT (k/g protein)	6.87 (5.54-7.38)	3.33 (2.39-3.84)	5.70 (3.88-6.95)	0.007	p ¹⁻² = 0.002

*p < 0.05 was considered statistically significant. The Kruskal–Wallis test was used. **p < 0.017 was considered statistically significant. Bonferroni correction has been made. Mann-Whitney U test was used.

MDA; malondialdehyde; SOD: superoxide dismutase; CAT: catalase.

Table 2. Histopathological assay; JTBS results

Groups	JTBS mean values Median (Q1-Q3)	p*	Post-hoc**
Group 1	9.5 (9.0-10.0)	< 0.001*	p ¹⁻² < 0.001 p ²⁻³ = 0.014
Group 2	3.0 (3.0-4.0)		
Group 3	8.0 (7.0-8.0)		

*p < 0.05 was considered statistically significant. Kruskal–Wallis test was used.

**p < 0.017 was considered statistically significant. Bonferroni correction has been made. Mann-Whitney U test was used.

JTBS: Johnsen testicular biopsy score.

testis, just as in other tissues, by disrupting the arterial circulation because the tunica of the testis is not elastic. Thrombi formed at venous, arterial, and capillary levels cause testicular ischemia and subsequent necrosis¹⁸. By eliminating the factor causing ischemia in reperfusion, blood flow to the ischemic organ can be restored, irreversible damage can be prevented, and lost functions can be restored. However, “reperfusion damage” occurs due to the free oxygen radicals formed and the activated blood cells in the blood. Reperfusion after ischemia damages the tissue more than ischemia^{19,20}.

Despite the search for alternative medical treatment methods for testicular torsion, emergency surgery is still a valid and widely applied treatment. Although the effect of testicular tissue by torsion is related to the duration and degree of torsion, the efficacy of numerous pharmacological agents and treatments have been evaluated in animal models to prevent testicular reperfusion injury since the primary preventable tissue damage occurs during the post-detorsion reperfusion period²¹.

Exosomes are serious cell-cell communication messengers and contain proteins, lipids, and genetic material that can be transported between cells to mediate intercellular communication and immune response²². Their existence as natural carriers, high biodistribution and stability in plasma, and relatively small size make them suitable candidates for therapeutic applications¹².

A recent study suggested that exosomes produced from human breast milk may reduce oxidative stress and support immune system development²³, and this encouraged us to use human breast milk exosomes in our study. Admyr et al. showed that exosomes in human breast milk can potentially affect the infant's immune system¹⁴. To investigate the effect of

exosomes against I/R damage in testicular torsion due to their protective effects in reperfusion injury mentioned above and positive effects on sperm motility and viability, we used exosomes obtained from breast milk, which has not been tested in the literature before.

First, we showed that the presence of human breast milk exosome produced in the Department of Biochemistry of KTU was confirmed by the presence of CD63 and the absence of calnexin using the Western blot method. Then we applied ip low dose (1 cc) exosomes 30 min before detorsion to the torsted testis. Studies show that local injection of the damaged testis has no adverse effect on the contralateral testis^{24,25}. Our study, in which we obtained statistically significant results, shows that testicular I/R damage and necrosis occur in the testicular tissue.

Keivan et al. have shown that rats' blood serum-derived exosomes inhibit testicular damage, including seminiferous tubule degeneration and loss of interstitial tissue, due to high NO and MDA levels following testicular torsion and NO, MDA, and apoptotic gene expression levels decrease and provide healthy sperm production²⁶. Similarly, in our study, when the MDA values of group 2 and group 3 were compared, we found a statistically significant difference between the two groups. Group 3 had significantly lower MDA values. The MDA value does not increase due to the effect of exosomes on reducing tissue damage. In addition, no statistically significant difference was found when the MDA values of group 1 and group 3 were compared.

No statistically significant difference was found when the SOD and CAT values of group 1 and group 3 were compared, and this makes us think that antioxidant enzymes are consumed less in group 3, and instead, the exosome is used as an antioxidant. As a result of these data, we think the exosome is biochemically effective in I/R damage since the MDA and SOD values are significantly different, and the CAT values are higher than Group 2, although there is no statistically significant difference.

Liu et al. showed that caspase-dependent apoptosis pathways are activated after testicular torsion injury, and spermatogenic cells are largely apoptotic, whereas treatment with exosomes significantly reduces the degree of apoptosis in spermatogenic cells²⁴. In addition, it has been shown histopathologically that the proliferative effect of the exosomes on spermatogenic cells is not limited to primary spermatocytes, and that it can inhibit spermatogenic cell

apoptosis and promote proliferation after testicular torsion-detorsion injury²⁴. In our study, when group 2 and group 3 were compared, it was shown that the testicles of group 3 had less damage than group 2, and the difference between them was statistically significant. When group 1 and group 3 were compared, although there were histopathological differences, there was no statistically significant difference. Since exosome application significantly reduced the reperfusion damage in testicular tissue compared to the torsion group in histopathology, it was seen as an expected result that it was not statistically significant. In group 3, spermatids and spermatozoa were preserved, although not as much as normal; it has been shown that there is a significant difference compared to group 2.

Finally, the fact that exosomes were taken from human breast milk, prepared and confirmed with a series of procedures, and the application dose was taken by taking samples from previous experimental studies can be considered among the limitations of our study.

Conclusion

Exosomes are important in diagnostic and therapeutic research as they are natural liposomes. Although studies on exosomes obtained from human breast milk have succeeded and increased in recent years, there is no clinical use in humans yet. We showed that exosomes were significantly protecting the testicular tissue and germ cells compared to the torsion group and new long-term experimental studies with different doses should be needed.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols, obtained informed consent from patients, and received approval from the Ethics Committee. The SAGER guidelines were followed according to the nature of the study.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

1. Yılmaz E, Hizli F, Afşarlar ÇE, Demirtaş C, Apaydin S, Karaman İ, et al. Early diagnosis of testicular torsion in rats by measuring plasma d-dimer levels: comparative study with epididymitis. *J Pediatr Surg*. 2015;50:651-4.
2. Mansour M, Degheili J, Khalifeh I, Tamim H, Jaafar RF, El-Hout Y. Remote ischemic conditioning in a rat model of testicular torsion: does it offer testicular protection? *J Pediatr Urol*. 2019;15:43.e1-7.
3. Akgür FM, Kiliç K, Aktuğ T. Reperfusion injury after detorsion of unilateral testicular torsion. *Urol Res*. 1993;21:395-9.
4. Asgari SA, Mokhtari G, Falahatkar S, Mansour-Ghanaei M, Roshani A, Zare A, et al. Diagnostic accuracy of C-reactive protein and erythrocyte sedimentation rate in patients with acute scrotum. *Urol J*. 2006;3:104-8.
5. Turner TT, Lysiak JJ. Oxidative stress: a common factor in testicular dysfunction. *J Androl*. 2008;29:488-98.
6. Shiraishi K, Naito K, Yoshida K. Nitric oxide promotes germ cell necrosis in the delayed phase after experimental testicular torsion of rat. *Biol Reprod*. 2001;65:514-21.
7. Cuzzocrea S, Riley DP, Caputi AP, Salvemini D. Antioxidant therapy: a new pharmacological approach in shock, inflammation, and ischemia/reperfusion injury. *Pharmacol Rev*. 2001;53:135-59.
8. Visser AJ, Heyns CF. Testicular function after torsion of the spermatic cord. *BJU Int*. 2003;92:200-3.
9. Miyake H, Lee C, Chusilp S, Bhalla M, Li B, Pitino M, et al. Human breast milk exosomes attenuate intestinal damage. *Pediatr Surg Int*. 2020;36:155-63.
10. Yıldırım C, Yuksek OH, Urkmez A, Sahin A, Somay A, Verit A. Protective effects of Tadalafil and darbepoetin against ischemia - reperfusion injury in a rat testicular torsion model. *Int Braz J Urol*. 2018;44:1005-13.
11. Becker EJ Jr., Turner TT. Endocrine and exocrine effects of testicular torsion in the prepubertal and adult rat. *J Androl*. 1995;16:342-51.
12. Kahraman T, Güçlüler G, Gürsel İ. Exosomes: natural nanovesicle candidates used in the diagnosis and treatment. *Türk J Immunol*. 2014;2:34-40.
13. Lässer C, Alikhani VS, Ekström K, Eldh M, Paredes PT, Bossios A, et al. Human saliva, plasma and breast milk exosomes contain RNA: uptake by macrophages. *J Transl Med*. 2011;9:9.
14. Admyre C, Johansson SM, Qazi KR, Filén JJ, Laheesmaa R, Norman M, et al. Exosomes with immune modulatory features are present in human breast milk. *J Immunol*. 2007;179:1969-78.
15. Sun Y, Oberley LW, Li Y. A simple method for clinical assay of superoxide dismutase. *Clin Chem*. 1988;34:497-500.
16. Aebi HE. Catalase. In: *Methods of Enzymatic Analysis*. Vol. 3. Weinheim: Verlag Chemie; 1987. p. 273-85.
17. Uchiyama M, Mihara M. Determination of malonaldehyde precursor in tissues by thiobarbituric acid test. *Anal Biochem*. 1978;86:271-8.
18. Bowlin PR, Gatti JM, Murphy JP. Pediatric testicular torsion. *Surg Clin North Am*. 2017;97:161-72.
19. Siemionow M, Arslan E. Ischemia/reperfusion injury: a review in relation to free tissue transfers. *Microsurgery*. 2004;24:468-75.
20. Lysiak JJ, Nguyen QA, Turner TT. Peptide and nonpeptide reactive oxygen scavengers provide partial rescue of the testis after torsion. *J Androl*. 2002;23:400-9.
21. Shimizu S, Tsounapi P, Dimitriadis F, Higashi Y, Shimizu T, Saito M. Testicular torsion-detorsion and potential therapeutic treatments: a possible role for ischemic postconditioning. *Int J Urol*. 2016;23:454-63.
22. Wang B, Zhong Y, Gao C, Li J. Myricetin ameliorates scopolamine-induced memory impairment in mice via inhibiting acetylcholinesterase and down-regulating brain iron. *Biochem Biophys Res Commun*. 2017;490:336-42.
23. Martin C, Patel M, Williams S, Arora H, Brawner K, Sims B. Human breast milk-derived exosomes attenuate cell death in intestinal epithelial cells. *Innate Immun*. 2018;24:278-84.
24. Liu H, Shi M, Li X, Lu W, Zhang M, Zhang T, et al. Adipose mesenchymal stromal cell-derived exosomes prevent testicular torsion injury via activating PI3K/AKT and MAPK/ERK1/2 pathways. *Oxid Med Cell Longev*. 2022;2022:8065771.
25. Hsiao CH, Ji AT, Chang CC, Cheng CJ, Lee LM, Ho JH. Local injection of mesenchymal stem cells protects testicular torsion-induced germ cell injury. *Stem Cell Res Ther*. 2015;6:113.
26. Keivan M, Mansouri Torghabeh F, Davoodi S, Moradi Maryamneghari S, Dadfar R. Single intratesticular injection of blood-serum-derived exosomes can potentially alleviate testopathy following testicular torsion. *Animal Model Exp Med*. 2022;5:362-8.